



US006316002B1

(12) **United States Patent**
Liu et al.

(10) Patent No.: **US 6,316,002 B1**
(45) Date of Patent: **Nov. 13, 2001**

(54) **GERMINATION ACTIVATED RED
GANODERMA LUCIDUM SPORES AND
METHOD FOR PRODUCING THE SAME**

(76) Inventors: Xin Liu, Building No. 391, 135
Xingang Xi Road, Guangzhou (CN);
Chee-Keung Chung, Room 1505,
Argyle Centre, 688 Nathan Rd.,
Mongkok, Kowloon (KR)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: 09/524,508

(22) Filed: Mar. 13, 2000

Related U.S. Application Data

(60) Provisional application No. 60/158,377, filed on Oct. 12,
1999.

(51) Int. Cl.⁷ A61K 35/00; A61K 35/78

(52) U.S. Cl. 424/195.15; 424/780

(58) Field of Search 424/195.15, 780

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,472,907 9/1984 Wada et al. 47/1.1
5,595,742 • 1/1997 Fujiwara et al. .

FOREIGN PATENT DOCUMENTS

1101860 • 4/1995 (CN).
1111529 • 11/1995 (CN).
1118005 • 3/1996 (CN).
1122658 • 5/1996 (CN).
1134306A • 10/1996 (CN).
1165032A • 11/1997 (CN).
1245718 • 8/1998 (CN).
1200286 • 12/1998 (CN).
1205231 • 1/1999 (CN).
1231925 • 10/1999 (CN).
2240026 9/1990 (JP).
52041208 8/1993 (JP).

OTHER PUBLICATIONS

Jiang et al. Shipin Kexue (Beijing). vol. 17, No. 4, pp.
19-22, CAPLUS Abstract enclosed, 1996.*
Yang et al. Zhongcaoyao. vol. 28, No. 12, pp. 721-723,
CAPLUS Abstract enclosed, 1997.*
Lee, Seung Y.; Cardiovascular Effects of Mycelium Extract
of *Ganoderma lucidum*: Inhibition of Sympathetic Outflow
as a Mechanism of Its Hypotensive Action, Chem Pharm.
Bull. vol. 38, p. 1359-1364 (1990).
Kim et al., Int. J. Mol. Med vol. 4(3), p. 273-277 (1999).
Lin et al., J. Ethnopharmacol., vol. 47 (1), p. 33-41 (1995).
Mekkawy, SaHar et al., Anti-HIV-1 and Anti-HIV-A-Pro-
tease Substances from *Ganoderma Lucidum*; *Phytochemis-
try*, vol. 49(6), p. 1651-1657 (1998).
Wasser, Solomon P., et al., Therapeutic Effects of Substances
Occurring in Higher Basidiomycetes Mushrooms: A Modern
Perspective; *Crit. Rev. Immunol.*, vol. 19(1), p. 65-96
(1999).

Miyazaki, Toshio et al., Studies on Fungal Polysacchar-
idesXXVII Structural Examination of a Water-soluble, Anti-
tumor Polysaccharide of *Ganoderma lucidum*; *Chem. Pharm.
Bull.*, vol. 29(12), p. 3611-3616 (1981).

Min, Byung-Sun et al.; Triterpenes from the Spores of
Ganoderma lucidum and Their Inhibitory Activity against
HIV-1 Protease; *Chem. Phar. Bull.*, vol. 46(10), p.
1607-1612 (1998).

Kino, K. et al., An immunomodulating protein, Ling
Zhi-8(LZ-8) prevents insulinitis an non-obese diabetic mice;
Diabetologia, vol. 33, p. 713-718 (1990).

Lieuwe, G. et al.; Ling Zhi-8: Studies of a New Immuno-
modulating Agent; *Transplantation*, vol. 60, p. 438-443
(1995).

Kino, Kohsuke et al.; Immunomodulator, LZ-8, Prevents
Antibody Production in Mice; *Int. J. Immunopharmac.*, vol.
13(8), p. 1109-1115 (1991).

Maruyama, Hirofumi; Antitumor Activity of *Sarcodon*
aspratus (BERK>)S. Ito and *Ganoderma lucidum* (FR.)
Karst.; *J. Pharmacobio-Dyn.*, vol. 12, p. 118-123 (1989).

Shimizu, Akria et al.; Isolation of an Inhibitor of Platelet
Aggregation from a Fungus, *Ganoderma lucidum*; *Chem.
Pharm. Bull.*, vol. 33, p. 3012-3015 (1985).

Morigiwa, Aiko et al.; Angiotension Converting
Enzyme-Inhibitory Triterpenes from *Ganoderma lucidum*;
Chem. Pharm. Bull., vol. 34(7), p. 3025-3028 (1986).

Kanmatsuse, Katsuo et al.; Studies of *Ganoderma lucidum*.
I. Efficacy against Hypertension and Side Effects; *Yakugaku
Zasshi*, vol. 105(10), p. 942-947 (1985).

Lieu, Chien-Whei et al.; The Effect of *Ganoderma lucidum*
on Induction of Differentiation in Leukemic U937 Cells;
Anticancer Research, vol. 12, p. 1211-1216 (1992).

(List continued on next page.)

Primary Examiner—Christopher R. Tate

(74) *Attorney, Agent, or Firm*—Fei-Fei Chao; Venable,
Baetjer, Howard & Civiletti, LLP

(57) **ABSTRACT**

The present invention describes a method for germination
activating spores of red *Ganoderma lucidum* to produce
bioactive substances which has medicinal effects on patients
with immunological disorders, cancer, AIDS, hepatitis,
diabetes, and cardiovascular diseases, and can prevent or
inhibit free radical oxidation and hepatotoxic effects. The
method can be subdivided into three stages. At the first stage,
a germination activation method is introduced which
includes soaking the spores in a solution to induce
germination, and placing the germination treated spores in a
culture box to induce the synthesis of bioactive substances
and softening of the cell walls of the spores. At the second
stage, sporoderm-broken ganoderma spores are collected by
treating the episporia with cell wall breaking enzymes
and/or mechanical force. At the last stage, the bioactive
substances are extracted from the sporoderm-broken spores
by drying at low temperature followed by extraction.

22 Claims, No Drawings